

Typical separating invariants

M. Domokos

Rényi Institute of Mathematics, Hungarian Academy of Sciences,
P.O. Box 127, 1364 Budapest, Hungary, E-mail: domokos@renyi.hu

Abstract

It is shown that a trivial version of polarization is sufficient to produce separating systems of polynomial invariants: if two points in the direct sum of the G -modules W and m copies of V can be separated by polynomial invariants, then they can be separated by invariants depending only on $\leq 2 \dim(V)$ variables of type V ; when G is reductive, invariants depending only on $\leq \dim(V) + 1$ variables suffice. Similar result is valid for rational invariants. Explicit bounds on the number of type V variables in a typical system of separating invariants are given for the binary polyhedral groups, and this is applied to the invariant theory of binary forms.

MSC: 13A50, 14L24

1 Introduction

Let G be a group, k an arbitrary base field, V, W finite dimensional G -modules over k . Write V^m for the m -fold direct sum $V \oplus \cdots \oplus V$. The coordinate ring $k[V]$ of V (the symmetric tensor algebra of the dual G -module V^*) contains the subalgebra $k[V]^G$ of polynomial invariants. We say that $v, v' \in V$ *can be separated (by polynomial invariants)* if there exists an $f \in k[V]^G$ with $f(v) \neq f(v')$. Following [4], Definition 2.3.8, we say that $S \subseteq k[V]^G$ is a *separating set* if whenever $v, v' \in V$ can be separated, then there exists an $f \in S$ with $f(v) \neq f(v')$.

We shall be interested in separating sets in $k[W \oplus V^m]^G$, where m is varying in $\mathbb{N}$. Given $f \in k[W \oplus V^d]$, $m \geq d \in \mathbb{N}$, and $1 \leq i_1 < \cdots < i_d \leq m$, denote by $f^{(i_1, \dots, i_d)}$ the function $f \circ \pi_{(i_1, \dots, i_d)} : W \oplus V^m \rightarrow k$, where $\pi_{(i_1, \dots, i_d)}$ is the projection mapping $(w, v_1, \dots, v_m) \mapsto (w, v_{i_1}, \dots, v_{i_d})$.

Definition 1 We say that $S \subseteq k[W \oplus V^d]^G$ is a complete system of typical separating invariants in type V variables relative to W , if for all $m \in \mathbb{N}$, $m \geq d$, we have that

$$S^{(m)} = \{f^{(i_1, \dots, i_d)} \mid f \in S, \quad 1 \leq i_1 < \dots < i_d \leq m\}$$

is a separating set in $k[W \oplus V^m]^G$. (Note that in this case for $q < d$, substituting zeros into the type V variables of index $q+1, q+2, \dots, d$ in the elements of S we obtain a separating set in $k[W \oplus V^q]^G$.)

This concept can be considered as a naive version of polarization, as the elements of $S^{(m)}$ are very special polarizations of the elements of S (see Section 7 for comments on polarization). Whereas the general polarization process is a basic tool (at least in characteristic zero) to produce generating invariants, the analogues of the above concept for generating systems seems to be useless. However, it turns out to be meaningful when we deal with separating sets.

Definition 2 Denote by $\sigma(G, W, V)$ the minimal non-negative integer d such that $k[W \oplus V^d]^G$ contains a complete system of typical separating invariants in type V variables relative to W .

In Section 2 we verify the general bound $\sigma(G, W, V) \leq 2 \dim(V)$. For a reductive group G this is improved in Section 3 to $\sigma(G, W, V) \leq 1 + \dim(V)$; the latter bound is sharp. We prove in Section 4 that the supremum of the numbers $\sigma(G, W, V)$ for a fixed finite group G can be interpreted as a purely group theoretic invariant that we call the Helly dimension of G . In Section 5 we turn to the invariant theory of binary forms: we investigate the numbers $\sigma(G, W, V)$ when G is the special linear group $\mathrm{SL}(2, \mathbb{C})$ and V is an irreducible $\mathrm{SL}(2, \mathbb{C})$ -module. It turns out that there is a universal constant upper bound for $\sigma(G, W, V)$, independent of W and V . In Section 6 we apply the idea of Definition 1 for rational invariants (where the general polarization method can not be applied).

2 General bound for the number of variables

Denote by $Gr(V \oplus V)$ the set of linear subspaces in $V \oplus V$.

Lemma 1 Whether $v = (w, v_1, \dots, v_m)$ and $v' = (w', v'_1, \dots, v'_m)$ can be separated or not depends only on the triple $(w, w', \mathrm{Span}_k\{v_1 \oplus v'_1, \dots, v_m \oplus v'_m\}) \in W \times W \times Gr(V \oplus V)$.

Proof. Take $v = (w, v_1, \dots, v_m) \in W \oplus V^m$, $v' = (w', v'_1, \dots, v'_m) \in W \oplus V^m$, $u = (w, u_1, \dots, u_d) \in W \oplus V^d$, $u' = (w', u'_1, \dots, u'_d) \in W \oplus V^d$ with

$$\mathrm{Span}_k(v_1 \oplus v'_1, \dots, v_m \oplus v'_m) = \mathrm{Span}_k(u_1 \oplus u'_1, \dots, u_d \oplus u'_d).$$

In particular, there are coefficients $\alpha_{ij} \in k$ such that

$$v_i \oplus v'_i = \sum_{j=1}^d \alpha_{ij}(u_j \oplus u'_j) \quad \text{for } i = 1, \dots, m.$$

Given an invariant $f \in k[W \oplus V^m]^G$, define $h \in k[W \oplus V^d]$ by

$$h(x, y_1, \dots, y_d) = f(x, \sum_{j=1}^d \alpha_{1j}y_j, \dots, \sum_{j=1}^d \alpha_{mj}y_j),$$

where x (resp. y_j) stand for vector variables of type W (resp. V). Obviously, h is G -invariant. By construction we have $h(u) = f(v)$ and $h(u') = f(v')$. So if $f(v) \neq f(v')$, then $h(u) \neq h(u')$. Consequently, if v and v' can be separated, then u and u' can be separated. The reverse implication is shown in the same way. $\square$

Theorem 2 *Let S be a separating set in $k[W \oplus V^{2n}]^G$, where $n = \dim(V)$. Then S is a complete system of typical separating invariants in type V variables relative to W . That is, we have $\sigma(G, W, V) \leq 2 \dim(V)$.*

Proof. Take a pair of points $v = (w, v_1, \dots, v_m)$, $v' = (w', v'_1, \dots, v'_m)$ in $W \oplus V^m$ where $m \geq 2n$. Since $\dim(\text{Span}_k\{v_1 \oplus v'_1, \dots, v_m \oplus v'_m\})$ is at most $2n$, there are indices $1 \leq i_1 < \dots < i_{2n} \leq m$ with

$$\text{Span}_k\{v_{i_1} \oplus v'_{i_1}, \dots, v_{i_{2n}} \oplus v'_{i_{2n}}\} = \text{Span}_k\{v_1 \oplus v'_1, \dots, v_m \oplus v'_m\}.$$

If v and v' can be separated, then $(w, v_{i_1}, \dots, v_{i_{2n}})$ and $(w', v'_{i_1}, \dots, v'_{i_{2n}})$ can be separated by Lemma 1, so there exists an $f \in S$ separating them. This means that $f^{(i_1, \dots, i_{2n})} \in k[W \oplus V^m]^G$ separates v and v' . $\square$

3 Reductive groups

Throughout this section we assume that k is algebraically closed. It is well known that if G is reductive (acting algebraically on V), then two points in V can be separated if and only if the Zariski closures of their G -orbits are disjoint, see for example Corollary 3.5.2 in [9]. Recall also that the Zariski closure of any G -orbit contains a unique closed orbit, hence a set of invariants is separating if it separates closed orbits.

Lemma 3 *Suppose that $v = (w, v_1, \dots, v_m) \in W \oplus V^m$, and for some $q \leq m$ we have that $\text{Span}_k\{v_1, \dots, v_q\} = \text{Span}_k\{v_1, \dots, v_m\}$. Then the G -orbit of v is closed in $W \oplus V^m$ if and only if the G -orbit of $(w, v_1, \dots, v_q)$ is closed in $W \oplus V^q$.*

Proof. There are coefficients $\beta_{q+i,j} \in k$ with $v_{q+i} = \sum_{j=1}^q \beta_{q+i,j} v_j$ holding for $i = 1, \dots, m-q$. The map

$$\begin{aligned} \varphi : W \oplus V^q &\rightarrow W \oplus V^m \\ (x, y_1, \dots, y_q) &\mapsto (x, y_1, \dots, y_q, \sum_{j=1}^q \beta_{q+1,j} y_j, \dots, \sum_{j=1}^q \beta_{m,j} y_j) \end{aligned}$$

is an injective morphism of affine G -varieties, its image is a G -stable linear subspace L of $W \oplus V^m$. So φ induces an isomorphism of the affine G -varieties $W \oplus V^q$ and L , moreover, φ maps the G -orbit of $(w, v_1, \dots, v_q)$ in $W \oplus V^q$ onto the G -orbit of v in $W \oplus V^m$. Hence the G -orbit of $(w, v_1, \dots, v_q)$ is closed in $W \oplus V^q$ if and only if the G -orbit of v is closed in L . Trivially, a subset of L is closed in L if and only if it is closed in $W \oplus V^m$. $\square$

Theorem 4 *Assume that G is reductive, and let S be a separating set in $k[W \oplus V^{n+1}]^G$, where $n = \dim(V)$. Then S is a complete system of typical separating invariants in type V variables relative to W . In other words, $\sigma(G, W, V) \leq \dim(V) + 1$ if G is reductive.*

Proof. Take $v = (w, v_1, \dots, v_m) \in W \oplus V^m$ and $v' = (w', v'_1, \dots, v'_m) \in W \oplus V^m$ where $m \geq n + 1$, and assume that the G -orbits of v and v' are different and closed with respect to the Zariski topology. We need to show that after a possible rearrangement of the type V variables, v and v' can be separated by an element of S (identifying $f \in S$ with $f^{(1, \dots, n+1)}$).

Without loss of generality we may assume that

$$\dim(\text{Span}_k\{v_1, \dots, v_m\}) \geq \dim(\text{Span}_k\{v'_1, \dots, v'_m\}),$$

and that $v_1, \dots, v_q$ is a basis of $\text{Span}_k\{v_1, \dots, v_m\}$ for some $q \leq n$.

Case I. $\text{Span}_k\{v'_1, \dots, v'_q\} \neq \text{Span}_k\{v'_1, \dots, v'_m\}$

Our assumptions imply that $v'_1, \dots, v'_q$ are linearly dependent, so $\lambda_1 v'_1 + \dots + \lambda_q v'_q = 0$ for some $\lambda_i \in k$, not all zero. It follows that the closure of the G -orbit of $(w', v'_1, \dots, v'_q)$ is contained in the proper linear subspace L of $W \oplus V^q$ defined by the system of linear equations $\sum_{i=1}^q \lambda_i y_i = 0$. On the other hand, the G -orbit of $(w, v_1, \dots, v_q)$ is not contained in L , and is closed in $W \oplus V^q$ by Lemma 3. Consequently, $(w, v_1, \dots, v_q)$ and $(w', v'_1, \dots, v'_q)$ can be separated by an element of $k[W \oplus V^q]^G$, implying that v and v' can be separated by an element of S .

Case II. $\text{Span}_k\{v'_1, \dots, v'_q\} = \text{Span}_k\{v'_1, \dots, v'_m\}$

By Lemma 3, the G -orbits of $(w, v_1, \dots, v_q)$ and $(w', v'_1, \dots, v'_q)$ are both closed in $W \oplus V^q$. If these G -orbits are different, then they can be separated by an element of $k[W \oplus V^q]^G$, whence they can be separated by an element of S . Otherwise replace v' by $g \cdot v'$ with a $g \in G$ satisfying $g \cdot v'_i = v_i$ for $i = 1, \dots, q$ and $g \cdot w' = w$. In other words, we may assume from

now on that $w = w'$, $v_1 = v'_1, \dots, v_q = v'_q$, and $v_1, \dots, v_q$ is a basis of $\text{Span}_k\{v_1, \dots, v_m\} = \text{Span}_k\{v'_1, \dots, v'_m\}$. There are unique coefficients $\gamma_{ij}, \gamma'_{ij} \in k$ such that

$$\sum_{j=1}^q \gamma_{ij} v_j = v_{q+i}, \quad \sum_{j=1}^q \gamma'_{ij} v'_j = v'_{q+i} \quad (i = 1, \dots, m - q).$$

Since v and v' are different, $v_{q+i} \neq v'_{q+i}$ for some i , say $v_{q+1} \neq v'_{q+1}$, implying that $\gamma_{1,j} \neq \gamma'_{1,j}$ for some $j \in \{1, \dots, q\}$. It follows that the G -orbits of $(w, v_1, \dots, v_{q+1})$ and $(w', v'_1, \dots, v'_{q+1})$ are different in $W \oplus V^{q+1}$ (since the action of G preserves the linear dependency relations among the type V variables). These orbits are closed in $W \oplus V^{q+1}$ by Lemma 3, hence they can be separated by an invariant in $k[W \oplus V^{q+1}]^G$. Now $q \leq n$, so this shows that v and v' (after a possible rearrangement of the type V variables) can be separated by an element of S . $\square$

Remark. Similar arguments imply that the nullcone is defined by invariants depending only on $\dim(V)$ variables of type V .

The following example shows that the bound $\dim(V) + 1$ on the number of type V variables in Theorem 4 is sharp.

Example. Let G be a torus $(k^\times)^{n-1}$ of rank $n - 1$ acting on $V = k^n$ by the rule

$$(\alpha_1, \dots, \alpha_{n-1}) \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} (\alpha_1 \cdots \alpha_{n-1})x_1 \\ \alpha_1^{-2}x_2 \\ \alpha_2^{-2}x_3 \\ \vdots \\ \alpha_{n-1}^{-2}x_n \end{bmatrix}.$$

Denote by $e_1, \dots, e_n$ the standard basis vectors in k^n , and set $v = (-e_1, e_1, e_2, e_3, \dots, e_n)$ and $v' = (e_1, e_1, e_2, e_3, \dots, e_n)$. Write $x(i)_j$ for the coordinate function on V^{n+1} mapping an $(n + 1)$ -tuple of vectors to the j th coordinate of the i th component. Then

$$x(1)_1 x(2)_1 x(3)_2 x(4)_3 x(5)_4 \cdots x(n+1)_n$$

is an invariant in $k[V^{n+1}]^G$ separating v and v' (provided that the characteristic of k is different from 2). On the other hand, we show that v and v' can not be separated by invariants depending only on $\leq n$ vector variables. Define the multidegree of a monomial in $k[V^{n+1}]$ to be $(d_1, \dots, d_n)$ if its total degree in the variables $x(1)_i, \dots, x(n+1)_i$ is d_i for $i = 1, \dots, n$. It is easy to see that an element of $k[V^{n+1}]$ is G -invariant if and only if all of its non-zero monomials have multidegree of the form $(2d, d, d, \dots, d)$ with $d \in \mathbb{N}$. Now let f be such a monomial. Then either $f(v) = 0 = f(v')$, or

$$f = x(1)_1^b x(2)_1^{2d-b} x(3)_2 x(4)_3 x(5)_4 \cdots x(n+1)_n^d \quad (b \in \{0, 1, \dots, 2d\}).$$

In the latter case, if f involves only n vector variables, then $b = 0$ or $b = 2d$, implying that $f(-e_1, u) = f(e_1, u)$ for all $u \in V^n$. If the characteristic of the base field is 2, then we modify the action to

$$(\alpha_1, \dots, \alpha_{n-1}) \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} (\alpha_1 \cdots \alpha_{n-1})x_1 \\ \alpha_1^{-3}x_2 \\ \alpha_2^{-3}x_3 \\ \vdots \\ \alpha_{n-1}^{-3}x_n \end{bmatrix},$$

and show in the same way as above that $(\omega e_1, e_1, e_2, \dots, e_n)$ (where ω is a primitive third root of 1) and $(e_1, e_1, e_2, \dots, e_n)$ can not be separated by invariants depending on $\leq n$ vector variables.

4 Finite groups

Throughout this section we assume that the group G is finite. Then all orbits are closed for any linear action of G , and two points in a G -module can be separated by polynomial invariants if and only if their orbits are different. This implies upper bounds for $\sigma(G, W, V)$ given in terms of the subgroup lattice of G .

Define the *Helly dimension* $\kappa(G)$ of G as the minimal natural number n with the following property: if we are given $m \geq n$ left cosets in G such that any n of them have a non-empty intersection, then the m cosets have a common element. We say that the subgroups $G_1, \dots, G_m$ are *intersection independent* if no G_i contains $\cap_{j \neq i} G_j$, the intersection of the others. Note that in this case $G_1, \dots, G_m$ are necessarily distinct, and if $m > 1$, then the G_i are proper subgroups. Denote by $\mu(G)$ the maximal size of an intersection independent set of subgroups of G , and write $\lambda(G)$ for the maximal length of a chain of proper subgroups in G .

Remark. The name "Helly dimension" is motivated by Helly's Theorem (cf. [7]). Numbers related to $\kappa(G)$ have already been introduced and studied for certain finite groups, see for example [3].

Lemma 5 *We have the inequalities $\kappa(G) \leq \mu(G) + 1$ and $\mu(G) \leq \lambda(G)$.*

Proof. Take an $n \geq \mu(G) + 1$ and $n + 1$ left cosets $g_0 G_0, g_1 G_1, \dots, g_n G_n$ in G such that any n of them have a non-empty intersection. Choose an element g from $\cap_{i=1}^n g_i G_i$. Since $G_1, \dots, G_n$ are not intersection independent, one of them, say G_n contains $\cap_{i=1}^{n-1} G_i$. By assumption there is an element h in $\cap_{i=0}^{n-1} g_i G_i = g_0 G_0 \cap g(\cap_{i=1}^{n-1} G_i)$, so h is contained also in $g G_n = g_n G_n$, showing that the intersection of all the cosets $g_i G_i$ ($i = 0, \dots, n$) is non-empty. This proves the first inequality.

For the second inequality observe that if $G_1, \dots, G_n$ is intersection independent, then $G_1 \supset G_1 \cap G_2 \supset G_1 \cap G_2 \cap G_3 \supset \dots \supset \cap_{i=1}^n G_i$ is a strictly descending chain of subgroups, and if $n > 1$, then G_1 is proper. $\square$

The Helly dimension is a very natural characteristic of a group. Given an action of G on some set X , $d \leq m \in \mathbb{N}$, consider the diagonal action on $X^m = X \times \dots \times X$, and ask whether two points $x = (x_1, \dots, x_m)$ and $x' = (x'_1, \dots, x'_m)$ belong to the same orbit, provided that $(x_{i_1}, \dots, x_{i_d})$ and $(x'_{i_1}, \dots, x'_{i_d})$ belong to the same orbit in X^d for all $1 \leq i_1 < \dots < i_d \leq m$. The minimal d such that this holds for all actions of G and all $m \geq d$ is nothing but $\kappa(G)$. The following variant of this observation explains our interest in $\kappa(G)$.

Proposition 6 *The Helly dimension $\kappa(G)$ is the maximal value of $\sigma(G, W, V)$, as W, V range over all finite dimensional G -modules.*

Proof. Recall that two points in a G -module can not be separated by polynomial invariants if and only if their orbit is the same.

First we prove the inequality $\sigma(G, W, V) \leq \kappa(G)$ for any G -modules W, V . Assume that $m \geq d = \kappa(G)$, $v = (w, v_1, \dots, v_m) \in W \oplus V^m$, $v' = (w', v'_1, \dots, v'_m) \in W \oplus V^m$, and suppose that for all $1 \leq i_1 < \dots < i_d \leq m$ we have that $\pi_{(i_1, \dots, i_d)}(v)$ and $\pi_{(i_1, \dots, i_d)}(v')$ belong to the same G -orbit in $W \oplus V^d$. We need to show that v and v' belong to the same G -orbit. Replacing v by an appropriate element in its orbit we may assume that $w = w'$. The set of elements g in the stabilizer of w with the property $gv_i = v'_i$ is a left coset gG_i in G , for $i = 1, \dots, m$. Our assumption is that any d of these cosets have a non-empty intersection. Since $d = \kappa(G)$, it follows that there is a common element g of these cosets, and obviously $g \cdot v = v'$.

To show the reverse inequality, take a natural number $d < \kappa(G)$. Then there is an $m > d$ and left cosets $g_1G_1, \dots, g_mG_m$ such that any d of them have a non-empty intersection, but the m cosets have no common element. It is well known that each subgroup of a finite group can be realized as a stabiliser of a vector in an appropriate finite dimensional representation (consider the representation induced by the trivial representation of the subgroup). Taking direct sums one can construct a finite dimensional G -module V and vectors $v_1, \dots, v_m \in V$ such that the stabilizer of v_i is G_i for $i = 1, \dots, m$. Set $v = (v_1, \dots, v_m) \in V^m$ and $v' = (g_1v_1, \dots, g_mv_m) \in V^m$. Our assumptions imply that $\pi_{(i_1, \dots, i_d)}(v)$ and $\pi_{(i_1, \dots, i_d)}(v')$ belong to the same G -orbit for all $1 \leq i_1 < \dots < i_d \leq m$, but v and v' belong to different orbits. This shows that $\sigma(G, 0, V) > d$. Since this holds for all $d < \kappa(G)$, we conclude that the supremum of the numbers $\sigma(G, W, V)$ as W and V vary is not smaller than $\kappa(G)$. $\square$

In the rest of this section we work out an explicit bound for the Helly dimension of the finite subgroups of $\text{SL}(2, \mathbb{C})$ (the binary polyhedral groups); this will be used in Section 5.

Recall that up to isomorphism the finite subgroups of SO_3 (the group of rotations of the 3-dimensional real Euclidean space) are the following: the cyclic group C_n of order n (the group of rotations of a regular n -angle pyramid), the dihedral group D_n of order $2n$ (the group of rotations stabilizing a regular n -gon), the alternating group A_4 (the group of rotations of a regular tetrahedron), the symmetric group S_4 (the group of rotations of a cube), and the alternating group A_5 (the group of rotations of a regular icosahedron). Denote by $\tilde{D}_n$, $\tilde{A}_4$, $\tilde{S}_4$, $\tilde{A}_5$ the preimages of these groups under the natural double covering $SU_2 \rightarrow SO_3$. Now up to isomorphism the following is a complete list of finite subgroups of $SL(2, \mathbb{C})$ (see for example Chapter 0.13 in [11]):

$$C_n, \quad \tilde{D}_n \quad (n = 1, 2, \dots), \quad \tilde{A}_4, \quad \tilde{S}_4, \quad \tilde{A}_5.$$

Proposition 7 *We have $\kappa(C_n) = 2$ for any $n > 1$.*

Proof. When $n > 1$, consider the cosets of two different elements with respect to the trivial subgroup $\{1\}$; this example shows that $\kappa(G) \geq 2$. To show the reverse inequality, denote by g a generator of C_n . Any coset in C_n is of the form $G_{q,r} = \{g^x \mid x \equiv r \pmod{q}\}$ for some positive divisor q of n and $r \in \{0, 1, \dots, q-1\}$. Take $m \geq 2$ and m cosets $G_{q_1, r_1}, \dots, G_{q_m, r_m}$ such that any pair has a non-empty intersection. Consider the system of congruences

$$x \equiv r_1 \pmod{q_1}; \quad x \equiv r_2 \pmod{q_2}; \quad \dots; \quad x \equiv r_m \pmod{q_m}.$$

By assumption, any pair of these congruences has a simultaneous solution. Therefore the system has a solution by the Chinese Remainder Theorem. This means that the intersection of the m given cosets is non-empty. $\square$

Remark. For comparison we mention that $\mu(C_n)$ can be arbitrary large: for example, if n is the product of r distinct primes $p_1, \dots, p_r$, then the subgroups of order $\frac{n}{p_1}, \frac{n}{p_2}, \dots, \frac{n}{p_r}$ are intersection independent.

Proposition 8 *We have $\kappa(\tilde{D}_n) \leq 4$ for any $n \geq 2$.*

Proof. In terms of generators and relations, $\tilde{D}_n = \langle a, b \mid b^4 = 1, \quad a^n = b^2, \quad bab^{-1} = a^{-1} \rangle$. For a positive divisor q of $2n$ and $r \in \{0, 1, \dots, q-1\}$, set $A_{q,r} = \{a^x \mid x \equiv r \pmod{q}\}$ and $B_{q,r} = \{ba^x \mid x \equiv r \pmod{q}\}$. The subgroups of $\tilde{D}_n$ are $A_{q,0}$, $A_{q,0} \cup B_{q,r}$, and the left cosets are $A_{q,r}$, $B_{q,r}$, $A_{q,r} \cup B_{q,s}$ (q a positive divisor of $2n$, $r, s \in \{0, 1, \dots, q-1\}$).

Take now $m+1 \geq 5$ cosets $g_i G_i$ ($i = 0, 1, \dots, m$) in $\tilde{D}_n$, and assume that any 4 of them have a common element. Partition each given coset as $g_i G_i = A^i \cup B^i$, where A^i is either empty or of the form $A_{q,r}$, and B^i is either empty or of the form $B_{q,r}$.

Case I. $A^i \cap A^j$ is non-empty for all $i < j$: Then by the proof of Proposition 7 we have that the intersection of all the A^i is non-empty, and we are done.

Case II. There two among the A^i , say A^0 and A^1 such that $A^0 \cap A^1 = \emptyset$. Then $g_0 G_0 \cap g_1 G_1 = B^0 \cap B^1$, hence for any $2 \leq i < j \leq m$ we have that $g_0 G_0 \cap g_1 G_1 \cap g_i G_i \cap g_j G_j = B^0 \cap B^1 \cap B^i \cap B^j$ is non-empty. Write q_j for the positive divisor of $2n$ and $r_j \in \{0, 1, \dots, q_j - 1\}$ with $B^j = B_{q_j, r_j}$, $j = 0, \dots, m$. Consider the following system of congruences:

$$x \equiv r_0 \pmod{q_0}; \quad x \equiv r_1 \pmod{q_1}; \quad x \equiv r_2 \pmod{q_2}; \quad \dots; \quad x \equiv r_m \pmod{q_m}.$$

Our assumption implies that for any $2 \leq i < j \leq m$, the subsystem consisting of the congruences with indices $0, 1, i, j$ has a solution. Consequently, any pair of the above $m + 1$ congruences has a simultaneous solution. Therefore the whole system has a solution by the Chinese Remainder Theorem. In other words, the $m + 1$ cosets have a non-empty intersection. $\square$

Corollary 9 *The Helly dimension of a finite subgroup of $\mathrm{SL}(2, \mathbb{C})$ is bounded by 6.*

Proof. An inspection of the corresponding subgroup lattices shows that $\lambda(A_4) = 3$, $\lambda(S_4) = 4$, and $\lambda(A_5) = 4$. It is easy to see that if $G \cong \tilde{G}/N$, where N is a two-element normal subgroup in the finite group $\tilde{G}$, then $\lambda(\tilde{G}) = \lambda(G) + 1$. Hence we have $\lambda(\tilde{A}_4) = 4$, $\lambda(\tilde{S}_4) = 5$, and $\lambda(\tilde{A}_5) = 5$. By Lemma 5 we get the desired bound for the Helly dimension of these groups. The two infinite series of finite subgroups of $\mathrm{SL}(2, \mathbb{C})$ are dealt with in Propositions 7 and 8. $\square$

5 Binary forms

Throughout this section our group G is the special linear group $\mathrm{SL}(2, \mathbb{C})$, $V = \mathrm{Pol}_d(\mathbb{C}^2)$ is the space of homogeneous binary forms of degree d with complex coefficients, endowed with the natural action of $\mathrm{SL}(2, \mathbb{C})$, whereas W is an arbitrary finite dimensional rational $\mathrm{SL}(2, \mathbb{C})$ -module. Recall that $\mathrm{Pol}_d(\mathbb{C}^2)$, $d = 0, 1, 2, \dots$ is a complete list of representatives of the isomorphism classes of irreducible rational G -modules. The study of the invariants of several binary forms is the most classical topic in invariant theory. Our aim here is to investigate the number of type V variables needed in a typical system of separating invariants in this case. It turns out that there is a uniform upper bound on $\sigma(G, W, V)$ (independent of W and V).

Theorem 10 *We have $\sigma(\mathrm{SL}(2, \mathbb{C}), W, V) \leq 8$ for any finite dimensional $\mathrm{SL}(2, \mathbb{C})$ -module W and irreducible $\mathrm{SL}(2, \mathbb{C})$ -module V ; that is, if S is a separating subset in $\mathbb{C}[W \oplus V^8]^{\mathrm{SL}(2, \mathbb{C})}$ (eight copies of V), then S is a typical system of separating invariants in type V variables relative to W .*

Remark. An arbitrary finite dimensional $\mathrm{SL}(2, \mathbb{C})$ -module is isomorphic to $V_1^{m_1} \oplus \cdots \oplus V_r^{m_r}$ (where $V_1, \dots, V_r$ are pairwise non-isomorphic irreducible $\mathrm{SL}(2, \mathbb{C})$ -modules). Theorem 10 implies that $\mathbb{C}[\bigoplus_{i=1}^r V_i^{m_i}]^{\mathrm{SL}(2, \mathbb{C})}$ contains a finite separating set whose elements depend only on ≤ 8 variables of type V_i for each $i = 1, \dots, r$.

As we mentioned already in Section 3, two points in a G -module can be separated by a polynomial invariant if and only if the unique closed orbits in their orbit closures are different. So Theorem 10 is equivalent to the following.

Proposition 11 *Let $v = (w, v_1, \dots, v_m)$ and $v' = (w', v'_1, \dots, v'_m)$ be points in $W \oplus V^m$ whose $\mathrm{SL}(2, \mathbb{C})$ -orbits are closed. Assume that for all $0 \leq r \leq 8$, $1 \leq i_1 < \cdots < i_r \leq m$, the closures of the orbits of $(w, v_{i_1}, \dots, v_{i_r})$ and $(w', v'_{i_1}, \dots, v'_{i_r})$ in $W \oplus V^r$ have a non-empty intersection. Then v and v' belong to the same orbit.*

In order to prove this statement we shall extensively use the generalization of the Hilbert-Mumford Criterion due to Birkes-Richardson (see [1] or Theorem 6.9 in [11]).

BRHM Criterion: Assume that the G -orbit of $u_1 \in W$ is closed. Then u_1 belongs to the Zariski closure of the G -orbit of u_2 if and only if there exists a *one-parameter subgroup* (shortly 1-PS) $\rho : \mathbb{C}^\times \rightarrow G$ (a homomorphism of algebraic groups) such that $\lim_{z \rightarrow 0} \rho(z)u_2$ exists and belongs to the G -orbit of u_1 .

Any non-trivial 1-PS in $\mathrm{SL}(2, \mathbb{C})$ is of the following form: there is an element $g \in \mathrm{SL}(2, \mathbb{C})$ and a natural number n such that

$$\rho(z) = g \begin{pmatrix} z^{-n} & 0 \\ 0 & z^n \end{pmatrix} g^{-1}. \quad (1)$$

We say that a 1-PS *acts on a linear form* $l \in \mathrm{Pol}_1(\mathbb{C}^2)$ *by a positive character* if the line $\mathbb{C}l$ is stabilized by ρ and $\rho(z)l = z^n l$ for a positive integer n . Note that if ρ is of the form (1), then the only lines fixed by ρ are spanned by gx and gy (where x and y stand for the usual coordinate functions on $\mathbb{C}^2$), and ρ acts by a positive character on gx (but not on gy). An orbit is called *maximal* if it is not contained in the closure of another orbit. We say that the linear form l is a *root of multiplicity* n of the binary form v if l^n divides v , and l^{n+1} does not divide v . From the BRHM Criterion and (1) one derives the following known facts.

Proposition 12 (i) *Let ρ be a 1-PS as in (1), and v a homogeneous binary form of degree d . Then $\lim_{z \rightarrow 0} \rho(z)v$ exists if and only if gx is a root of multiplicity $\geq d/2$ of v . If the multiplicity is greater than $d/2$, then $\lim_{z \rightarrow 0} \rho(z)v = 0$, and if d is even and the multiplicity is $d/2$, then $\lim_{z \rightarrow 0} \rho(z)v$ is a non-zero scalar multiple of $(gx)^{d/2}(gy)^{d/2}$.*

(ii) *If $\lim_{z \rightarrow 0} \rho(z)v$ exists for some non-trivial 1-PS, then it does not belong to the $\mathrm{SL}(2, \mathbb{C})$ -orbit of v , unless $v = 0$ or d is even and $v = (l_1 l_2)^{d/2}$ for some independent linear forms l_1, l_2 .*

Proposition 13 *The $\mathrm{SL}(2, \mathbb{C})$ -orbit of $v = (w, v_1, \dots, v_m) \in W \oplus V^m$ is closed and maximal if the components $v_1, \dots, v_m$ have no common root of multiplicity $\geq d/2$ (in all v_i). In particular, the orbit of v is closed and maximal if there is a type V component that has no root of multiplicity $\geq d/2$, or if there are two type V components that have no common root of multiplicity $\geq d/2$.*

Proof. Assume that the orbit of v is not closed. By the BRHM Criterion, there is a non-trivial 1-PS ρ such that $\lim_{z \rightarrow 0} \rho(z)v$ exists. Then $\lim_{z \rightarrow 0} \rho(z)v_i$ exists for all i . We may assume that ρ is of the form (1). Then gx is a root of all v_i with multiplicity $\geq d/2$ by Proposition 12. Next, assume that the orbit of v is not maximal. Then there is some $u \in W \oplus V^m$ and a non-trivial 1-PS ρ (of the form (1)) with $v = \lim_{z \rightarrow 0} \rho(z)u$, implying that $v_i = \lim_{z \rightarrow 0} \rho(z)u_i$ for all i . If d is odd, then all the v_i are zero, and if d is even, then all $v_i \in \mathbb{C}(gx)^{d/2}(gy)^{d/2}$ by Proposition 12, hence say gx is a common root of the v_i with multiplicity $d/2$. $\square$

Similar arguments yield the following consequence of Proposition 12.

Proposition 14 *Assume that the orbit of $v = (w, v_1, \dots, v_m) \in W \oplus V^m$ is closed. Then it is maximal if either of the following conditions holds:*

- (i) *v has a type V component which is non-zero, and can not be written as $(l_1 l_2)^{d/2}$ where d is even and l_1, l_2 are independent linear forms;*
- (ii) *v has two linearly independent components of type V .*

Remark. A description of the orbit closures is given in [8] for irreducible $\mathrm{SL}(2, \mathbb{C})$ -modules and in [10] for arbitrary $\mathrm{SL}(2, \mathbb{C})$ -modules. One could deduce Propositions 13 and 14 from the results of [10].

We shall need also the following elementary fact on the irreducible representations of $\mathrm{SL}(2, \mathbb{C})$.

Lemma 15 *The pointwise stabilizer of a linear subspace of dimension ≥ 2 in the irreducible $\mathrm{SL}(2, \mathbb{C})$ -module V is finite.*

Proof. Let H be the pointwise stabilizer in $\mathrm{SL}(2, \mathbb{C})$ of a subspace of V of dimension ≥ 2 . Any element of H is conjugate in $\mathrm{SL}(2, \mathbb{C})$ to one of $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ or $\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}$ or $\begin{pmatrix} z & 0 \\ 0 & z^{-1} \end{pmatrix}$ ($z \in \mathbb{C}^\times$). Direct computation shows that the fixed point subspaces of the unipotent elements are 1-dimensional. Therefore any element of H is semisimple. It follows that the connected component of the identity in H is contained in a maximal torus T of $\mathrm{SL}(2, \mathbb{C})$. Now T is

1-dimensional (as an algebraic variety), and the fixed point subspace of T is 1-dimensional if d is even and 0-dimensional if d is odd. Consequently, the connected component of the identity in H is 0-dimensional, hence is the trivial subgroup $\{1\}$, showing that H is finite. $\square$

Proof of Proposition 11. If all v_i, v'_i are zero, then the assertion holds trivially. From now on we assume that this is not the case.

Case I. There is a type V component of v , say v_1 , that has no root of multiplicity $\geq d/2$.

We claim that for all $0 \leq r \leq 7$ and $2 \leq i_1 < \dots < i_r \leq m$, the G -orbits of $(w, v_1, v_{i_1}, \dots, v_{i_r})$ and $(w', v'_1, v'_{i_1}, \dots, v'_{i_r})$ coincide. Indeed, the closures of these orbits intersect by assumption. On the other hand, the first G -orbit is closed and maximal by Proposition 13. These two facts clearly imply that the orbits coincide.

Case I/a. $v_i = \alpha_i v_1$ with $\alpha_i \in \mathbb{C}$ for $i = 2, \dots, m$.

Since (w, v_1, v_i) and (w', v'_1, v'_i) belong to the same G -orbit, we have also $v'_i = \alpha_i v'_1$ for $i = 2, \dots, m$. Take $g \in G$ with $g(w, v_1) = (w', v'_1)$, such a g exists, and $gv_i = g(\alpha_i v_1) = \alpha_i gv_1 = \alpha_i v'_1 = v'_i$ for $i = 2, \dots, m$, so $gv = v'$.

Case I/b. There is an $i \geq 2$, say $i = 2$ such that v_1 and v_2 are linearly independent.

Our assumptions imply that $u = (w, v_1, v_2)$ and $u' = (w', v'_1, v'_2)$ belong to the same G -orbit, so replacing v by an appropriate element in its orbit we may assume that $u = u'$. By the assumptions, for all $1 \leq r \leq 6$ and $3 \leq i_1 < \dots < i_r \leq m$, the elements $(u, v_{i_1}, \dots, v_{i_r})$ and $(u, v'_{i_1}, \dots, v'_{i_r})$ belong to the same G -orbit, hence $(v_{i_1}, \dots, v_{i_r})$ and $(v'_{i_1}, \dots, v'_{i_r})$ belong to the same orbit with respect to the stabilizer H of u . Now H is a finite subgroup of $\mathrm{SL}(2, \mathbb{C})$ by Lemma 15, so we can conclude from Corollary 9 that $(v_3, \dots, v_m)$ and $(v'_3, \dots, v'_m)$ belong to the same H -orbit, or in other words, that v and v' belong to the same G -orbit.

Case II. All components of v of type V have a root of multiplicity $\geq d/2$, and there are two components, say v_1 and v_2 that have no common root of multiplicity $\geq d/2$ (both in v_1 and v_2).

The orbit of $(w, v_1, v_2, v_{i_1}, \dots, v_{i_r})$ is closed and maximal by Proposition 13, hence coincides with the orbit of $(w', v'_1, v'_2, v'_{i_1}, \dots, v'_{i_r})$ for all $0 \leq r \leq 6$ and $3 \leq i_1 < \dots < i_r \leq m$. Moreover, v_1 and v_2 are linearly independent. One concludes that the orbits of v and v' are the same just like in Case I/b.

Case III. There is a linear form l such that l^{e+1} divides all of $v_1, \dots, v_m$, where $d = 2e$ or $d = 2e + 1$, and not all v_i are zero.

Note that l is the only root of multiplicity $\geq d/2$ for each non-zero component v_i . Without loss of generality we may assume that v_1 is non-zero. By Proposition 12, $\lim_{z \rightarrow 0} \rho(z)(v_1, \dots, v_m)$ exists if and only if ρ acts by a positive character on l , and in this case $\lim_{z \rightarrow 0} \rho(z)(v_1, \dots, v_m)$ does not belong to the orbit of $(v_1, \dots, v_m)$. Hence in this case $\lim_{z \rightarrow 0} \rho(z)w$ does not exist (recall that the orbit of v is closed by assumption). Consequently, $\lim_{z \rightarrow 0} \rho(z)(w, v_1)$ does not exist for any non-trivial 1-PS, implying that the orbit of $(w, v_1, v_{i_1}, \dots, v_{i_r})$ is closed for all $r \geq 0$ and $2 \leq i_1 < \dots < i_r \leq m$. Moreover, these orbits are maximal by Proposition 14. Thus the assumptions of the Proposition imply that $(w, v_1, v_{i_1}, \dots, v_{i_r})$ and $(w', v'_1, v'_{i_1}, \dots, v'_{i_r})$ belong to the same orbit provided that $r \leq 7$. One finishes in the same way as in Case I.

Since the roles of v and v' are symmetric, one can deal with Cases I', II', III' (obtained by replacing v by v') in the same way as with Cases I, II, III.

Note that the proof is already complete for odd d . There are however further cases to consider when d is even, which we assume from now on. From now on we automatically assume that we are not in the cases covered so far. In particular, this implies that any two type V components of v have a common root of multiplicity $\geq d/2$, and the same holds for v' .

Case IV. There is a non-zero type V component of v or v' , say v_1 , which can not be written as $(l_1 l_2)^{d/2}$ with two independent linear forms l_1 and l_2 .

There is a unique (up to scalar) root l of multiplicity $\geq d/2$ of v_1 , and $l^{d/2}$ necessarily divides all the v_i . Since the orbit of v is closed, just like in Case III, we conclude from Propositions 12 and 14 that the orbit of $(w, v_1, v_{i_1}, \dots, v_{i_r})$ is closed and maximal for all $0 \leq r$ and $2 \leq i_1 < \dots < i_r \leq m$. One finishes in the same way as in Case I (or III).

From now on we assume also that we are not in Case IV. This means in particular that each non-zero type V component of v and v' can be written as the product of the $(d/2)$ th powers of two independent linear forms.

Case V. There is a non-zero linear form l and linear forms $l_1, \dots, l_m$ such that $v_i = l^{d/2} l_i^{d/2}$ for $i = 1, \dots, m$, and there are two among the l_i , say l_1 and l_2 that are linearly independent.

For any 1-PS ρ acting on l by a positive character $\lim_{z \rightarrow 0} \rho(z)(v_1, \dots, v_m)$ exists, furthermore, $\lim_{z \rightarrow 0} \rho(z)(v_1, v_2)$ does not belong to the orbit of (v_1, v_2) . Since the orbit of v is closed, this implies $\lim_{z \rightarrow 0} \rho(z)w$ does not exist for such ρ . Taking into account that l is the only common root of v_1 and v_2 , this implies that the orbit of $(w, v_1, v_2, v_{i_1}, \dots, v_{i_r})$ is closed for any $r \geq 0$ and $3 \leq i_1 < \dots < i_r \leq m$. These orbits are maximal by Proposition 14. One can finish as in Case I/b.

Case V'. Replace v by v' in Case V.

Case VI. $v_i = \alpha_i(l_1 l_2)^{d/2}$, $v'_i = \beta_i(b_1 b_2)^{d/2}$, $i = 1, \dots, m$, where $\alpha_i, \beta_i \in \mathbb{C}$, the linear forms l_1, l_2 are independent, and the linear forms b_1, b_2 are independent.

There is a non-zero scalar among the α_i, β_j , say $\alpha_1 \neq 0$. Then the G -orbit of (w, v_1) in $W \oplus V$ is closed by Lemma 3. If $v'_1 = 0$, then (w, v_1) is not contained in the closure of the orbit of (w', v'_1) , contradicting our assumption in the Proposition. So $\beta_1 \neq 0$, and the orbit of (w', v'_1) is closed. Moreover, the orbits of $(w, v_1, v_{i_1}, \dots, v_{i_r})$ and $(w', v'_1, v'_{i_1}, \dots, v'_{i_r})$ are also closed by Lemma 3 for all $r \geq 0$ and $2 \leq i_1 < \dots < i_r \leq m$, hence they coincide by our assumptions if $r \leq 7$. One finishes in the same way as in Case I/a.

For the rest of the proof we assume additionally that we are not in Cases V, V', VI. So $v_1, \dots, v_m$ have no common root of multiplicity $\geq d/2$. On the other hand, any pair v_i, v_j have a common root of multiplicity $\geq d/2$. Moreover, the same holds for v' . This implies that there are three pairwise independent linear forms l_1, l_2, l_3 such that a non-zero scalar multiple of each of $(l_1 l_2)^{d/2}$, $(l_1 l_3)^{d/2}$, $(l_2 l_3)^{d/2}$ occurs among the components, say $v_1 = (l_1 l_2)^{d/2}$, $v_2 = (l_1 l_3)^{d/2}$, $v_3 = (l_2 l_3)^{d/2}$, and all the v_i belong to $\mathbb{C}v_1 \cup \mathbb{C}v_2 \cup \mathbb{C}v_3$. By Proposition 13, the orbit of (w, v_1, v_2, v_3) is closed and maximal, hence by assumption coincides with the orbit of (w', v'_1, v'_2, v'_3) , and so we may assume that they are equal (replace v by an appropriate element in its orbit). It follows that all type V components of v' belong to $\mathbb{C}v_1 \cup \mathbb{C}v_2 \cup \mathbb{C}v_3$. Take a non-zero component v_i of v , where $i \geq 4$. Then there is a nonzero $\alpha_i \in \mathbb{C}$ and $f(i) \in \{1, 2, 3\}$ such that $v_i = \alpha_i v_{f(i)}$. Assume for example that $f(i) = 3$. Then the same argument as above shows that (w, v_1, v_2, v_i) and $(w', v'_1, v'_2, v'_i) = (w, v_1, v_2, v'_i)$ belong to the same orbit. Therefore $v'_i \notin \mathbb{C}v_1 \cup \mathbb{C}v_2$, so $v'_i = \beta_i v_3$ with a non-zero $\beta_i \in \mathbb{C}$. By assumption, the orbit closures of (v_3, v_i) and (v'_3, v'_i) have a non-empty intersection, implying that $\alpha_i = \beta_i$, hence $v_i = v'_i$. By symmetry in v and v' , we have that v_i is non-zero if and only if v'_i is non-zero. So a repeated use of the above argument implies that $v_i = v'_i$ for all $i \geq 4$, hence $v = v'$. $\square$

We take a digression and state and prove the analogues of Theorem 10 for ordinary polarization.

Theorem 16 *If S is a separating set of invariants in $\mathbb{C}[W \oplus V^3]^{\text{SL}(2, \mathbb{C})}$ (three copies of V), then the polarizations with respect to the type V variables of the elements of S constitute a separating set in $\mathbb{C}[W \oplus V^m]^{\text{SL}(2, \mathbb{C})}$ for an arbitrary m . Moreover, it is sufficient to consider polarizations of the following special form: when $m \geq 3$, take the coefficients of $f(w, v_{i_1}, v_{i_2}, \sum_{j=3}^{\min\{m, 8\}} t^{j-3} v_{i_j})$ (viewed as a polynomial in t), as f ranges over S , and $i_1, i_2, i_3, \dots$ are different elements of $\{1, \dots, m\}$; when $m < 3$, take $f(w, v_1, \dots, v_m, 0, \dots, 0)$ as f ranges over S .*

Proof. The proof is almost the same as the proof of Theorem 10. Write q for $\min\{m, 8\}$. We treat the same Cases as in that proof. The only difference is that when we reduced a given Case to the study of orbits of a finite group H (the stabilizer of (w, v_{i_1}, v_{i_2})), then we

need to point out that if $(v_{i_3}, \dots, v_{i_q})$ and $(v'_{i_3}, \dots, v'_{i_q})$ belong to different H -orbits, then there is a $t \in \mathbb{C}$ such that $\sum_{j=3}^q t^{j-3} v_{i_j}$ belongs to a different H -orbit than $\sum_{j=3}^q t^{j-3} v'_{i_j}$; this follows from the results on "cheap polarization" in [6]. Hence the closed maximal orbit of $(w, v_{i_1}, v_{i_2}, \sum_{j=3}^q t^{j-3} v_{i_j})$ is not contained in the closure of the orbit of $(w', v'_{i_1}, v'_{i_2}, \sum_{j=3}^q t^{j-3} v'_{i_j})$, implying that $f(w, v_{i_1}, v_{i_2}, \sum_{j=3}^q t^{j-3} v_{i_j}) \neq f(w', v'_{i_1}, v'_{i_2}, \sum_{j=3}^q t^{j-3} v'_{i_j})$ for an appropriate $f \in S$. $\square$

Remark. The bound 3 on the number of type V variables is sharp, as it can be seen already when $\dim(V) = 3$.

6 Rational invariants

Throughout this section k is algebraically closed. Our results on typical systems of separating invariants have a version for rational invariants as well. Given an irreducible G -variety X , denote by $k(X)$ the field of rational functions on X and by $k(X)^G$ its subfield of invariants. Following [11] we say that $R \subset k(X)^G$ *separates orbits in general position* if there is a non-empty open subset X_0 of X such that if $x, x' \in X_0$ belong to different orbits, then there is a rational invariant $f \in R$ defined both at x and x' such that $f(x) \neq f(x')$. It is well known that a finite set R of rational invariants separates orbits in general position if and only if $k(X)^G$ is a purely inseparable extension of its subfield generated by k and R ; see [11] for the characteristic zero case, and combine it with (18.2) from [2] to get the general statement. A theorem of Rosenlicht [12] asserts that a finite set R of rational invariants separating orbits in general position always exists. (Note that if a finite set R of rational invariants separates orbits in general position, then there is a non-empty G -stable open subset U in X such that the elements of R are all defined on U , and separate inequivalent points in U , thus a geometric quotient U/G exists.)

Write X^m for the product $X \times \dots \times X$ of m copies of X .

Theorem 17 *Let X be an irreducible G -subvariety of $V = k^n$ (where G acts linearly on V), let Y be an arbitrary irreducible G -variety, and assume that $R \subset k(Y \times X^{n+1})^G$ separates orbits in general position in $Y \times X^{n+1}$. Then for any $m \geq n + 1$ we have that*

$$R^{(m)} = \{f^{(i_1, \dots, i_{n+1})} \mid 1 \leq i_1 < \dots < i_{n+1} \leq m\}$$

separates orbits in general position in $Y \times X^m$. In particular, if $\text{char}(k) = 0$, then $R^{(m)}$ generates the field of rational invariants on $Y \times X^m$.

Proof. Let U be a non-empty open subset of $Y \times X^{n+1}$ such that inequivalent points in U can be separated by an element of R . The complement of U in $Y \times X^{n+1}$ is a proper closed

subset Z in $Y \times X^{n+1}$. For $1 \leq i_1 < \dots < i_{n+1} \leq m$, denote by $\pi_{(i_1, \dots, i_{n+1})}$ the projection morphism $Y \times X^m \rightarrow Y \times X^{n+1}$, $(y, x_1, \dots, x_m) \mapsto (y, x_{i_1}, \dots, x_{i_{n+1}})$. Then

$$Z^{(m)} = \bigcup_{1 \leq i_1 < \dots < i_{n+1} \leq m} \pi_{(i_1, \dots, i_{n+1})}^{-1}(Z)$$

is a proper closed subset of $Y \times X^m$; denote its non-empty complement by $U^{(m)}$. (If U was G -stable, then $U^{(m)}$ is G -stable, since the projection maps are G -equivariant.) Note that if $x \in U^{(m)}$, then any projection $\pi_{(i_1, \dots, i_{n+1})}(x)$ belongs to U . If for $x, x' \in U^{(m)}$ there is an $(i_1, \dots, i_{n+1})$ such that $y = \pi_{(i_1, \dots, i_{n+1})}(x)$ and $y' = \pi_{(i_1, \dots, i_{n+1})}(x')$ belong to different G -orbits, then y and y' can be separated by some $f \in R$, and so $f^{(i_1, \dots, i_{n+1})}$ separate x and x' .

Assume now that all projections $\pi_{(i_1, \dots, i_{n+1})}$ map $x = (y, x_1, \dots, x_m) \in U^{(m)}$ and $x' = (y', x'_1, \dots, x'_m)$ into the same G -orbit in $Y \times X^{n+1}$. After a possible renumbering of the indices, we may assume that $x_1, \dots, x_n$ span the k -subspace $\text{Span}_k\{x_1, \dots, x_m\}$ of V . So there are coefficients α_{ij} with $x_{n+i} = \sum_{j=1}^n \alpha_{ij} x_j$ for $i = 1, \dots, m - n$. Since $(y, x_1, \dots, x_n, x_{n+i})$ and $(y', x'_1, \dots, x'_n, x'_{n+i})$ belong to the same G -orbit of $Y \times X^{n+1}$ for all i by our assumption, we have that $x'_{n+i} = \sum_{j=1}^n \alpha_{ij} x'_j$ holds for $i = 1, \dots, m - n$ (recall that the action of G on X is induced by a linear action on V). It follows that for an arbitrary element $g \in G$ with $g \cdot (y, x_1, \dots, x_n) = (y', x'_1, \dots, x'_n)$ we have that $g \cdot x = x'$. Such a g exists by our assumption, hence x and x' belong to the same G -orbit of $Y \times X^m$. $\square$

7 Concluding remarks

The present paper is motivated by Weyl's Theorem [13], Theorem 2.5A, asserting that if k is of characteristic zero, then $k[W \oplus V^m]^G$ is generated by the polarizations (with respect to the type V variables) of the elements of a generating system of $k[W \oplus V^n]^G$, where $n = \dim(V)$. It is well known that this does not hold when $\text{char}(k)$ is positive (see for example [5] for a class of examples). However, in a recent preprint, J. Draisma, G. Kemper and D. Wehlau [6] proved that polarizing the elements of a separating set in $k[W \oplus V^n]^G$, $n = \dim(V)$, one always gets a separating set in $k[W \oplus V^m]^G$, regardless of the characteristic of the base field.

Our statements Theorem 2 and 4 are weaker if one is interested in degree bounds for separating invariants. On the other hand, they are "really" about separating invariants, in the sense that they have no analogues for generating invariants, even in characteristic zero. From the point of view of polarizations, our results can be viewed as an addendum to [6] making possible some computational simplifications in the construction of a separating set in $k[W \oplus V^m]^G$ for large m . Namely, one takes a separating set S_0 in $k[W \oplus V^n]^G$ with $n = \dim(V)$. Then it is sufficient to take those polarizations of the elements of S_0 that depend only on $\leq 2n$ (resp. $\leq n + 1$ for reductive G) type V variables to get a (multi)homogeneous separating set in $k[W \oplus V^m]^G$. This is an essential simplification from the point of view of explicit computations.

Finally, this trivial version of "polarization" applies also for rational invariants, where the general polarization process does not make sense.

Acknowledgement

I thank P. P. Pálffy for remarks that led to an improved presentation of Section 4. This research was partially supported by OTKA No. T046378, the Bolyai Fellowship and Leverhulme Research Exchange Grant F/00158/X.

References

- [1] D. Birkes, Orbits of linear algebraic groups, *Ann. Math. II.*, Ser. 93 (1971), 459-475.
- [2] A. Borel, *Linear algebraic groups*, Second edition, Graduate Texts in Mathematics, 126. Springer-Verlag, New York, 1991.
- [3] P. J. Cameron and Ph. Cara, Independent generating sets and geometries for symmetric groups, *J. Algebra* 258 (2002), 641-650.
- [4] H. Derksen and G. Kemper, *Computational Invariant Theory*, Springer-Verlag, Berlin, 2002.
- [5] M. Domokos, Matrix invariants and the failure of Weyl's theorem, in "Polynomial Identities and Combinatorial Methods" (ed. A. Giambruno, A. Regev, M. Zaicsev), *Lecture Notes in Pure and Applied Mathematics* 235 (2003), 215-236, Marcel Dekker.
- [6] J. Draisma, G. Kemper and D. Wehlau, Polarization of separating invariants, preprint (2005).
- [7] Ed. Helly, Über Mengen konvexen Körper mit gemeinschaftlichen Punkten, (German), *J. Deutsche Math.*, Ver 32 (1923), 175-176.
- [8] D. Khadziev, *Invariant Theory of Binary Forms*, Tashkent: Branch of the Academy of Sciences, 52 pp. (Russian), 1978.
- [9] P. E. Newstead, *Introduction to Moduli Problems and Orbit Spaces*, Springer-Verlag, Berlin, 1978.
- [10] V. L. Popov, The structure of orbit closures in finite dimensional linear representations of $SL(2)$, (Russian), *Matem. Zametki* 16 (1974), 943-950.
- [11] V. L. Popov and E. B. Vinberg, Invariant theory, in: *Algebraic Geometry IV*, *Encyclopaedia of Mathematical Sciences* 55, Springer-Verlag, Berlin, Heidelberg, 1994.

- [12] M. Rosenlicht, A remark on quotient spaces, An. Acad. Brasil. Ci. 35 (1963), 487-489.
- [13] H. Weyl, The Classical Groups, Princeton University Press, Princeton, 1939.